

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
17 October 2002 (17.10.2002)

PCT

(10) International Publication Number
WO 02/080754 A2

(51) International Patent Classification⁷: **A61B**
(21) International Application Number: PCT/US02/10238
(22) International Filing Date: 3 April 2002 (03.04.2002)
(25) Filing Language: English
(26) Publication Language: English
(30) Priority Data:
60/281,277 3 April 2001 (03.04.2001) US
(71) Applicant (*for all designated States except US*): **THE-
SEUS IMAGING CORPORATION** [US/US]; 101 Arch
Street, Suite 1762, Boston, MA 02110 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— *without international search report and to be republished upon receipt of that report*

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(72) Inventor; and
(75) Inventor/Applicant (*for US only*): **GREEN, Allan, M.** [US/US]; 19 Francis Avenue, Cambridge, MA 02138 (US).
(74) Agents: **DECONTI, Giulio, A. Jr.** et al.; Lahive & Cockfield, LLP, 28 State Street, Boston, MA 02109 (US).

(54) Title: METHODS FOR USING ANNEXIN FOR DETECTING CELL DEATH IN VIVO AND TREATING ASSOCIATED CONDITIONS

(57) Abstract: The present invention provides methods and compositions for imaging cell death *in vivo*, as well as methods and compositions for tumor radiotherapy.



WO 02/080754 A2

METHODS FOR USING ANNEXIN FOR DETECTING CELL DEATH IN VIVO AND TREATING ASSOCIATED CONDITIONS

5

Related Applications

This application claims priority to U.S. Provisional Patent Application Serial No. 60/281,277 filed April 3, 2001, the entire contents of which are incorporated herein by reference.

10

Background of the Invention

Apoptosis and Necrosis

Apoptosis refers to "programmed cell death" whereby the cell executes a "cell suicide" program. It is now thought that the apoptosis program is evolutionarily conserved among virtually all multicellular organisms, as well as among all the cells in a particular organism. Further, it is believed that in many cases, apoptosis may be a "default" program that must be actively inhibited in healthy surviving cells.

The decision by a cell to submit to apoptosis may be influenced by a variety of regulatory stimuli and environmental factors (Thompson, 1995). Physiological activators of apoptosis include tumor necrosis factor (TNF), Fas ligand, transforming growth factor β , the neurotransmitters glutamate, dopamine, N-methyl-D-aspartate, withdrawal of growth factors, loss of matrix attachment, calcium and glucocorticoids. Damage-related inducers of apoptosis include heat shock, viral infection, bacterial toxins, the oncogenes *myc*, *rel* and *E1A*, tumor suppressor p53, cytolytic T-cells, oxidants, free radicals and nutrient deprivation (antimetabolites). Therapy-associated apoptosis inducers include gamma radiation, UV radiation and a variety of chemotherapeutic drugs, including cisplatin, doxorubicin, bleomycin, cytosine arabinoside, nitrogen mustard, methotrexate and vincristine. Toxin-related inducers or

apoptosis include ethanol and d-amyloid peptide.

Apoptosis can have particularly devastating consequences when it occurs pathologically in cells that do not normally regenerate, such as neurons. Because such cells are not replaced when they die, their loss can lead to debilitating and sometimes fatal dysfunction of the affected organ. Such dysfunction is evidenced in a number of neurodegenerative disorders that have been associated with increased apoptosis, including Alzheimer's disease, Parkinson's disease, amyotrophic lateral sclerosis, retinitis pigmentosa and cerebellar degeneration.

The consequences of undesired apoptosis can be similarly devastating in other pathologies as well, including ischemic injury, such as typically occurs in cases of myocardial infarction, reperfusion injury and stroke. In particular, apoptosis is believed to play a central role in very delayed infarction after mild focal ischemia (Du, *et al.*, 5 1996). Additional diseases associated with increased apoptosis include, but are not limited to, the following: AIDS; myelodysplastic syndromes, such as aplastic anemia; and toxin induced liver disease, including damage due to excessive alcohol consumption.

Necrosis is the localized death of cells or tissue due to causes other than 10 apoptosis (*i.e.*, other than the execution of the cell's intrinsic suicide program). Necrosis can be caused by traumatic injury, bacterial infection, acute hypoxia and the like. There is some overlap between the two types of cell death, in that some stimuli can cause either necrosis or apoptosis or some of both, depending on the severity of the injury.

15 Asymmetry of Biological Membranes

It is generally believed that biological membranes are asymmetric with respect to specific membrane phospholipids. In particular, the outer leaflet of eukaryotic plasma membranes is formed predominantly with the cholinephospholipids, such as sphingomyelin and phosphatidylcholine (PC), whereas the inner leaflet contains 20 predominantly aminophospholipids, such as phosphatidylserine (PS) and phosphatidylethanolamine (PE). This asymmetry is thought to be maintained by the activity of an adenosine triphosphate (ATP)-dependent aminophospholipid translocase, which selectively transports PS and PE between bilayer leaflets (Seigneuret and Devaux, 1984). Other enzymes thought to be involved in the transport of phospholipids between 25 leaflets include ATP-dependent floppase (Connor, *et al.*, 1992) and lipid scramblase (Zwaal, *et al.*, 1993).

Although asymmetry appears to be the rule for normal cells, the loss of such asymmetry is associated with certain physiological, as well as pathogenic, processes. For example, it has been recognized that membrane asymmetry, detected as appearance 30 of PS on the outer leaflet of the plasma membrane ("PS exposure"), is one of the earliest manifestations of apoptosis, preceding DNA fragmentation, plasma membrane blebbing, and loss of membrane integrity (Martin, *et al.*, 1995; Fadok, *et al.*, 1992).

Similar re-orientation has been observed in sickle cell disease (Lane, *et al.*, 1994) "B-thalassemia (Borenstain-Ben Yashar, *et al.*, 1993), platelet activation, and in 35 some mutant tumor cell lines with defective PS transport. A gradual appearance of PS on the outer leaflet has also been observed to occur in aging red blood cells (Tait and

Gibson, 1994). When the PS exposure on such cells reaches a threshold level, the cells are removed from circulation by macrophages (Pak and Fidler, 1991). All of the above conditions proximately culminate in the death of the affected cells (*i.e.*, cells with significant PS exposure).

5 It will be appreciated that PS exposure is a component in both apoptosis and necrosis. Its role in the initial stages of apoptosis is summarized above. Once the apoptotic cell has reached the terminal stages of apoptosis (*i.e.*, loss of membrane integrity), it will be appreciated that the PS in both plasma membrane leaflets will be "exposed" to the extracellular milieu. A similar situation exists in cell death by necrosis,
10 where the loss of membrane integrity is either the initiating factor or occurs early in the necrotic cell death process; accordingly, such necrotic cells also have "exposed" PS, since both plasma membrane leaflets are "exposed".

Annexin

15 Annexin V is normally found in high levels in the cytoplasm of a number of cells including placenta, Lymphocytes, monocytes, biliary and renal (cortical) tubular epithelium. Although the physiological function of annexins has not been fully elucidated, several properties of annexins make them useful as diagnostic and/or therapeutic agents. In particular, it has been discovered that annexins possess a very
20 high affinity for anionic phospholipid surfaces, such as a membrane leaflet having an exposed surface of phosphatidylserine (PS).

Summary of the Invention

 The present invention provides methods and compositions for imaging cell death
25 *in vivo*, as well as methods and compositions for tumor radiotherapy and phototherapy. The present invention is based, at least in part, on the discovery that the combination of an annexin with a contrast agent allows for the efficient and effective detection of cells undergoing cell death using magnetic resonance imaging. The present invention is also based, at least in part, on the discovery that the combination of an annexin with an
30 optically active molecule, such as a fluorescent dye, allows for the efficient and effective detection of cells undergoing cell death by optical imaging. Finally, the present invention is based, at least in part, on the discovery that administering a composition comprising an annexin coupled with a therapeutic radioisotope to a tumor bearing subject that has been treated with chemotherapeutic agent, allows for the specific and
35 enhanced delivery of the radiation carried by the annexin- therapeutic radioisotope composition to the tumor site.

Accordingly, the present invention provides a magnetic resonance imaging composition which includes an annexin, *e.g.*, annexin V, coupled to a contrast agent, such as a paramagnetic agent (*e.g.*, a gadolinium-chelating group complex, such as gadolinium-diethylenetriamine penta-acetic acid, or a lanthanum chelating group
5 complex) or a superparamagnetic agent (*e.g.*, a metal oxide, such as Fe, Co, Ni, Cu, Zn, As, Se, Mo, Tc, Ru, Rh, Pd, Ag, Cd, In, Sn, Re, Os, Ir, Pt, Au, Hg, Tl, Pb, Bi, Po, or At oxide). The metal oxide is preferably coated with a polymer, *e.g.*, dextran or variants thereof. The annexin may be coupled to the contrast agent directly or indirectly.

In another aspect, the present invention provides compositions comprising an
10 annexin, *e.g.*, annexin V, coupled to a contrast agent, such as a polymer coated metal oxide, and a radioisotope, *e.g.*, a diagnostic or therapeutic radioisotope. Such compositions are suitable for both MRI and nuclear medicine imaging. For example, the composition may include annexin V coupled to a contrast agent and a radioisotope (linked to the annexin via hydrazino nicotinamide (HYNIC)). In one embodiment, the
15 annexin may be coupled to a carrier that is cleared or metabolized by a desirable route. Examples of such carriers include, but are not limited to, dextran particles or colloidal particles or metal oxide particles, such as superparamagnetic iron oxide particles (which are typically phagocytosed in the liver).

In another aspect, the present invention provides a method for the *in vivo*
20 imaging of cell death, *e.g.*, cell death caused by apoptosis, in a mammalian subject, for example, in an organ of a mammalian subject or a portion thereof (*e.g.*, brain, heart, liver, lung, pancreas, colon) or a gland of a mammalian subject or a portion thereof (*e.g.*, prostate or mammary gland). The method includes administering to the subject a magnetic resonance imaging composition comprising annexin coupled to a contrast
25 agent; and obtaining a magnetic resonance image, wherein said image is a representation of cell death in the mammalian subject. In one embodiment, the magnetic resonance image is obtained 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 105, 110, 115, or 120 minutes after the administration of the magnetic resonance imaging composition to the subject. In another embodiment, the magnetic
30 resonance image is obtained about 12-30, 15-25, 20-25, or 20-30 hours after the administration of the magnetic resonance imaging composition to the subject. Ranges intermediate to the above recited values are also intended to be part of this invention. For example, ranges using a combination of any of the above recited values as upper and/or lower limits are intended to be included. In a preferred embodiment, the
35 magnetic resonance image is obtained at a plurality of time points, thereby monitoring

changes in the number of cells undergoing cell death or monitoring changes in the location of cells undergoing cell death.

The magnetic resonance imaging composition may be administered at a concentration of 1-1000 µg protein/kg, 1-900 µg protein/kg, 1-800 µg protein/kg, 1-700 µg protein/kg, 1-600 µg protein/kg, 1-500 µg protein/kg, 1-400 µg protein/kg, 1-300 µg protein/kg, 1-200 µg protein/kg, 1-100 µg protein/kg, 1-50 µg protein/kg, or 1-20 µg protein/kg. In another embodiment, the magnetic resonance imaging composition is administered intravenously, intraperitoneally, intrathecally, intrapleurally, intralymphatically, or intramuscularly.

10 In a further aspect, the present invention provides an optical imaging composition which includes an annexin, e.g., annexin V, coupled to a biologically compatible and optically active molecule, such as a fluorescent dye like fluorescein, which can be visualized during optical evaluations such as endoscopy, bronchoscopy, peritonoscopy, direct visualization, surgical microscopy and retinoscopy. Moreover, by
15 the appropriate choice of optically active molecule, an annexin-optically active molecule combination may be useful in photodynamic therapy (PDT), a novel approach for the treatment of cancer and other diseases, such as macular degeneration, which may be used as a primary or adjunctive therapeutic modality. In the present invention, PDT works by exposing an annexin molecule linked to a photosensitizing drug to specific
20 wavelengths of light in the presence of oxygen. When this reaction occurs, the normally innocuous photosensitizing molecule becomes cytotoxic via an activated species of oxygen, known as "singlet oxygen." The ability of annexin to localize at sites of tumor cell apoptosis makes this an ideal drug to use in combination with anti-cancer treatment which leads to apoptosis or necrosis of tumor cells. The temporal introduction of the
25 annexin-photosensitizing drug after induction of tumor cell apoptosis or necrosis creates a circumstance for differential localization of the annexin-photosensitizing molecule combination at the tumor site, providing the opportunity for additional tumor cell killing using appropriate light exposure. Typically, laser energy, delivered to the diseased tissue, e.g., cancer site, directly or through a fiberoptic device, chemically activates the
30 drug and creates a toxic form of oxygen which destroys the cancerous cells with minimal damage to healthy cells. Examples of optically active agents which could be used in PDT when linked to annexin include PHOTOFRIN®, Lutrin, ANTRIN®, FOSCAN®, aminolevulinic acid, aluminum (III) phthalocyanine tetrasulfonate, Hypericin, verteporfin, and methylene blue dye. Among the possible targets for PDT
35 are tumors of the brain, head and neck, breast, esophagus, lung, pleural cavity, ovary, abdominal cavity, bladder, prostate, cervix, skin, peritoneal cavity, eye and

aerodigestive system.

In yet another aspect, the present invention provides a method for imaging cell death in a mammalian subject *in vivo* by administering to the subject an optical imaging composition comprising annexin coupled to an optically active molecule; illuminating
5 the subject with a light source; and visually monitoring the presence of the optical imaging composition in the subject, thereby obtaining an image, wherein the image is a representation of cell death in the mammalian subject.

In another aspect, the present invention provides a composition comprising an annexin, *e.g.*, annexin V, coupled with a therapeutic radioisotope, *e.g.*, ^{103}Pd , ^{186}Re , ^{188}Re , ^{90}Y , ^{153}Sm , ^{159}Gd , or ^{166}Ho . The therapeutic radioisotope and the annexin may be
10 coupled at a ratio of 1:1, 1.1:1, 1.2:1, 1.3:1, 1.4:1, 1.5:1, 1.6:1, 1.7:1, 1.8:1, 1.9:1, or 2:1 (therapeutic radioisotope:annexin). Ranges of values using a combination of any of the above recited values as upper and/or lower limits are intended to be included in the present invention.

In a further aspect, the present invention provides a method of tumor radiotherapy by administering to a mammalian subject having a tumor an effective tumor reducing amount of a composition comprising an annexin coupled with a therapeutic radioisotope. The foregoing method may be used in conjunction with total body irradiation or targeted external irradiation and/or a treatment employing at least
20 one chemotherapeutic agent (*e.g.*, dimethyl busulfan, cyclophosphamide, bischloroethyl nitrosourea, cytosine arabinoside, or 6-thioguanine). In addition, the method may be used in conjunction with biologically active anti-cancer agents and apoptosis inducing agents such as TNF, TRAIL or Fas or with antibodies, small molecules or pharmacophores which bind these receptors and also induce apoptosis.

In another aspect, the present invention features a method of tumor radiotherapy, which includes treating a subject having a tumor with a chemotherapeutic agent and subsequently administering to the subject an effective tumor reducing amount of a composition comprising an annexin coupled with a therapeutic radioisotope.

The timing of the administration of the annexin coupled with a therapeutic agent
30 is critical to the effectiveness of the therapeutic intervention. The modified annexin should be administered at a time which assures its bioavailability at times of apoptosis or necrosis of the target tissue. Diagnostic imaging studies using radiolabeled annexin V indicate that the administration of a therapeutically modified annexin preferably should be within 24 hours of the completion of a course of chemotherapy of
35 lymphoma with multiple antimetabolite drugs (so-called, CHOP or MOPP therapy) to optimize the availability of annexin localization in the damaged tumor. Optimal time of

administration may be within 72 hours of chemotherapeutic treatment of solid tumors such as breast cancer, lung cancer or sarcoma as shown by imaging studies in patients. Use of a diagnostic imaging agent, such as radiolabeled annexin, to determine the extent of apoptosis may be used to qualify patients for administration of therapeutically modified annexin and to determine the optimal dose of therapeutically modified annexin.

Other features and advantages of the invention will be apparent from the following detailed description, and from the claims.

Brief Description of the Drawings

Figure 1 depicts the attachment of Annexin V to an iron oxide coated with the non-polymer DMSA.

Figure 2 depicts the attachment of Annexin V to a polymer coated magnetic iron oxide.

Detailed Description of the Invention

The present invention provides methods and compositions for imaging cell death *in vivo*, as well as methods and compositions for tumor radiotherapy. The present invention is based, at least in part, on the discovery that the combination of an annexin with a contrast agent allows for the efficient and effective detection of cells undergoing cell death using magnetic resonance imaging. The present invention is also based, at least in part, on the discovery that the combination of an annexin with an optically active molecule, such as a fluorescent dye, allows for the efficient and effective detection of cells undergoing cell death by optical imaging.

Finally, the present invention is based, at least in part, on the discovery that administering a composition comprising an annexin coupled with a therapeutic radioisotope to a tumor bearing subject that has been treated with chemotherapeutic agent, allows for the specific and enhanced delivery of the radiation carried by the annexin- therapeutic radioisotope composition to the tumor site. Without intending to be limited by mechanism, it is believed that the administration of the chemotherapeutic agent will cause apoptosis or necrosis at the site of the tumor, thereby allowing the annexin-therapeutic radioisotope complex to be specifically targeted to the site of the tumor and delivering the radiation, which will kill the cell to which the annexin binds, as well as neighboring cells.

Accordingly, the present invention provides a magnetic resonance imaging composition which includes an annexin, *e.g.*, annexin V, coupled (either directly or indirectly) to a contrast agent.

As used herein, a "contrast agent" is intended to include any agent that is physiologically tolerable and capable of providing enhanced contrast for magnetic resonance imaging. Contrast agents typically have the capability of altering the response of a tissue to magnetic fields. Contrast agents include paramagnetic agents, *e.g.*, a gadolinium-chelating group complex, such as gadolinium-diethylenetriamine penta-acetic acid, or a manganese chelating group complex; or biologically compatible superparamagnetic agents such as iron oxide. Contrast agents, such as those described in U.S. Patent No. 4,687,658; U.S. Patent No. 5,314,680; and U.S. Patent No. 4,976,950 are intended to be used in preparing the compositions of the present invention. Contrast agents are commercially available (*e.g.*, the gadolinium chelate ProhanceTM is available from Squibb and the gadolinium chelate DotaremTM is available from Guerbet).

A suitable contrast agent must preferably be biocompatible, *e.g.*, non-toxic, chemically stable, not absorbed by the body or reactive with a tissue, and eliminated from the body within a short time. In one embodiment, the contrast agent may be coupled to a carrier that is cleared or metabolized by a desirable route. Examples of such carriers include, but are not limited to, dextran particles or colloidal particles (which are typically phagocytosed in the liver).

In another aspect, the present invention provides a method for the *in vivo* imaging of cell death, *e.g.*, cell death caused by apoptosis, in a mammalian subject. The method includes administering to the subject a magnetic resonance imaging composition comprising annexin coupled to a contrast agent; and obtaining a magnetic resonance image, wherein said image is a representation of cell death in the mammalian subject.

As used herein, the term "cell death" includes the processes by which mammalian cells die. Such processes include apoptosis (both reversible and irreversible) and processes thought to involve apoptosis (*e.g.*, cell senescence), as well as necrosis. "Cell death" is used herein to refer to the death or imminent death of nucleated cells (*e.g.*, neurons, myocytes, hepatocytes and the like) as well as to the death or imminent death of anucleate cells (*e.g.*, red blood cells, platelets, and the like). Cell death is typically manifested by the exposure of PS on the outer leaflet of the plasma membrane.

As used herein, the term "subject" includes warm-blooded animals, preferably mammals, including humans. In a preferred embodiment, the subject is a primate. In an even more preferred embodiment, the subject is a human. Cell death may be imaged or detected in, for example, an organ of a subject or a portion or specimen thereof (*e.g.*,
5 brain, heart, liver lung, pancreas, colon) or a gland of a subject or a portion thereof (*e.g.*, prostate, pituitary or mammary gland). For example, cell death may be imaged or detected using surgical or needle biopsy of a subject after administration of the annexin to the subject; or by the use of a catheter that may detect radiation in a vessel of a subject.

10 As used herein, the term "administering" to a subject includes dispensing, delivering or applying a composition of the invention to a subject by any suitable route for delivery of the composition to the desired location in the subject, including delivery by either the parenteral or oral route, intramuscular injection, subcutaneous/intradermal injection, intravenous injection, buccal administration, transdermal delivery and
15 administration by the rectal, colonic, vaginal, intranasal or respiratory tract route.

The compositions of the invention may be administered to a subject in an amount effective, at dosages and for periods of time necessary, to achieve the desired result. An effective amount of the compositions of the invention may vary according to factors such as disease *state*, *e.g.*, the tumor stage, age, and weight of the subject, and the ability
20 of the composition to elicit a desired response in the subject. An effective amount is also one in which any toxic or detrimental effects (*e.g.*, side effects) of the compositions are outweighed by the therapeutically or diagnostically beneficial effects. The compositions of the invention may be administered at a concentration of, for example, 1-1000 µg protein/kg, 1-900 µg protein/kg, 1-800 µg protein/kg, 1-700 µg protein/kg, 1-
25 600 µg protein/kg, 1-500 µg protein/kg, 1-400 µg protein/kg, 1-300 µg protein/kg, 1-200 µg protein/kg, 1-100 µg protein/kg, 10-100 µg protein/kg, 10-80 µg protein/kg, 10-60 µg protein/kg, 10-40 µg protein/kg, or 10-20 µg protein/kg.

The magnetic resonance image may be obtained using any of the art known techniques, for example, using a Picker Corp. Whole Body Superconducting System
30 operating at 0.3 T using a 30 cm transmitter coil tuned to 0.26 T (10.08 MHz) or other MRI devices with field strengths ranging from 0.05 Tesla to 4.0 Tesla. Typically, the subject is placed in a powerful, highly uniform, static magnetic field. Magnetized protons (hydrogen nuclei) within the subject align like small magnets in this field. Radiofrequency pulses are then utilized to create an oscillating magnetic field
35 perpendicular to the main field, from which the nuclei absorb energy and move out of alignment with the static field, in a state of excitation. As the nuclei return from

excitation to the equilibrium state, a signal induced in the receiver coil of the instrument by the nuclear magnetization can then be transformed by a series of algorithms into images. Images based on different tissue characteristics can be obtained by varying the number and sequence of pulsed radiofrequency fields in order to take advantage of
5 magnetic relaxation properties of the tissues.

If it is desired to follow the localization and/or the signal over time, for example, to record the effects of a treatment on the distribution and/or localization of cell death, the imaging can be repeated at selected time intervals to construct a series of images. The intervals can be as short as minutes, or as long as days, weeks, months or years.
10 Images generated by methods of the present invention may be analyzed by a variety of methods. They range from a simple visual examination, mental evaluation and/or printing of a hardcopy, to sophisticated digital image analysis.

The magnetic resonance image may be obtained 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 105, 110, 115, or 120 minutes after the
15 administration of the magnetic resonance imaging composition to the subject. In another embodiment, the magnetic resonance image is obtained about 10-30, 15-25, 20-25, or 20-30 hours after the administration of the magnetic resonance imaging composition to the subject. In a preferred embodiment, the magnetic resonance image is obtained at a plurality of time points, thereby monitoring changes in the number of
20 cells undergoing cell death or monitoring changes in the location of cells undergoing cell death.

The present invention also provides an optical imaging composition which includes an annexin, *e.g.*, annexin V, coupled to an optically active molecule.

As used herein, an "optically active molecule" includes any molecule that has the
25 ability to be optically detected, for example, by the use of medically available visualization devices such as endoscopes, bronchoscopes and minimally invasive surgical devices using optical detection of anatomic structures. Examples of optically detectable molecules include fluorescein and methylene blue. Optically active molecules may also include those agents useful in photodynamic therapy (PDT). PDT
30 works by exposing an annexin molecule linked to a photosensitizing molecule to specific wavelengths of light in the presence of oxygen. When this reaction occurs, the normally innocuous photosensitizing drug becomes cytotoxic via an activated species of oxygen, known as "singlet oxygen." Examples of optically active agents which could be used in PDT when linked to annexin include PHOTOFRIN®, Lutrin, ANTRIN®,
35 FOSCAN®, aminolevulinic acid, aluminum (III) phthalocyanine tetrasulfonate, Hypericin, verteporfin, and methylene blue dye.

In another aspect, the present invention provides a composition comprising an annexin, *e.g.*, annexin V, coupled with a therapeutic radioisotope. As used herein, a "therapeutic radioisotope" is a radioisotope that is recognized as being useful and suitable for injection into a patient for therapeutic applications. A therapeutic radioisotope, as used herein, is preferably an alpha (α) or beta (β) emitting radioisotope. In one embodiment, the therapeutic radioisotope is not a gamma (γ) emitting radioisotope. Examples of therapeutic radioisotopes include ^{103}Pd , ^{186}Re , ^{188}Re , ^{90}Y , ^{153}Sm , ^{159}Gd , or ^{166}Ho , ^{131}I , ^{123}I , ^{126}I , ^{133}I , ^{111}In , and ^{113}In .

The therapeutic radioisotope and the annexin may be coupled at a ratio of 1:1, 1.1:1, 1.2:1, 1.3:1, 1.4:1, 1.5:1, 1.6:1, 1.7:1, 1.8:1, 1.9:1, or 2:1 (therapeutic radioisotope:annexin). Ranges of values using a combination of any of the above recited values as upper and/or lower limits are intended to be included in the present invention.

In a further aspect, the present invention provides a method of tumor radiotherapy by administering to a mammalian subject having a tumor an effective tumor reducing amount of a composition comprising an annexin coupled with a therapeutic radioisotope. The foregoing method may be used in conjunction with total body irradiation or targeted external irradiation and/or a treatment employing at least one chemotherapeutic agent (*e.g.*, dimethyl busulfan, cyclophosphamide, bischloroethyl nitrosourea, cytosine arabinoside, or 6-thioguanine). The appropriate timing for the administration of the annexin-therapeutic isotope composition may be determined using any of the imaging techniques described herein or the imaging techniques described in U.S. Patent No. 6,197,278 B1, the contents of which are incorporated herein by reference.

Various aspects of the invention are described in further detail in the following subsections:

Synthesis of Annexin Containing Compounds of the Invention

The invention can be practiced using purified native, recombinant, or synthetically-prepared annexin. Annexin V, for example, may be conveniently purified from human placenta (as described in Funakoshi, *et al.* (1987) *Biochemistry* 26:5572, the contents of which are incorporated herein by reference). Recombinant annexin offers several advantages, however, including ease of preparation and economic efficiency. A number of different annexins have been cloned from humans and other organisms. Their sequences are available in sequence databases, including GenBank.

The invention is preferably practiced using annexin V, for several reasons. First, annexin V is one of the most abundant annexins, (ii) it is simple to produce from natural or recombinant sources, and (iii) it has a high affinity for phospholipid membranes.

Human annexin V has a molecular weight of 36 kd and a high affinity ($K_d = 7 \text{ nmol/L}$) for phosphatidylserine (PS). The sequence of human annexin V can be obtained from GenBank under accession numbers U05760-U05770.

An exemplary expression system suitable for making annexin for use with the present invention employs the pET12a expression vector (Novagen, Madison, Wisconsin) in *E. coli*. (described in Wood, *et al.* (1996) *Blood* 88:1873-1880, incorporated herein by reference).

Other bacterial expression vectors may be utilized as well. They include, *e.g.*, the plasmid pGEX (Smith, *et al.* (1988) *Gene* 67:31) and its derivatives (*e.g.*, the pGEX series from Pharmacia Biotech, Piscataway, NJ). These vectors express the polypeptide sequences of a cloned insert fused in-frame with glutathione-S-transferase.

Recombinant pGEX plasmids can be transformed into appropriate strains of *E. coli* and fusion protein production can be induced by the addition of IPTG (isopropyl-thio galactopyranoside). Solubilized recombinant fusion protein can then be purified from cell lysates of the induced cultures using glutathione agarose affinity chromatography according to standard methods (described in, for example, Ausubel, *et al.* *Current Protocols in Molecular Biology* (John Wiley and Sons, Inc., Media, PA). Other commercially-available expression systems include yeast expression systems, such as the *Pichia* expression kit from Invitrogen (San Diego, CA); baculovirus expression systems (Reilly, *et al.* in *Baculovirus Expression Vectors: A Laboratory Manual* (1992); Clontech, Palo Alto CA); and mammalian cell expression systems (Clontech, Palo Alto CA; Gibco-BRL, Gaithersburg MD).

A number of features can be engineered into the expression vectors, such as leader sequences which promote the secretion of the expressed sequences into culture medium. The recombinantly produced polypeptides are typically isolated from lysed cells or culture media.

Isolated recombinant polypeptides produced as described above may be purified by standard protein purification procedures, including differential precipitation, molecular sieve chromatography, ion exchange chromatography, isoelectric focusing, gel electrophoresis and affinity chromatography. Protein preparations can also be concentrated by, for example, filtration (Amicon, Danvers, Mass.).

Annexin produced as described above may then be coupled to a contrast agent, an optically active molecule, or a therapeutic radioisotope. The particular contrast agent, optically active molecule, or therapeutic radioisotope selected will depend on the particular application the skilled artisan intends to use.

5

Annexin Radiolabeling

Annexins may be radiolabeled by a variety of methods known in the art (*e.g.*, as described in U.S. Patent No. 5,985,240; U.S. Patent No. 4,361,544 and U.S. Patent No. 4,427,646, the entire contents of each of which are incorporated herein by reference).

10 Annexins may be directly radioiodinated, through electrophilic substitution at reactive aromatic amino acids. Iodination may also be accomplished via pre-labeled reagents, in which the reagent is iodinated and purified, and then linked to the annexin. Iodination may also be achieved through the use of chelates, *e.g.*, DTPA and EDTA chelates, as described in, for example, U.S. Patent No. 4,986,979; U.S. Patent No. 4,479,930 and
15 U.S. Patent No. 4,668,503.

In selecting a suitable therapeutic radioisotope, the skilled artisan will typically consider factors including, but not limited to, (i) minimum of particle emission, (ii) primary photon energy of between about 50 and 500 keV, (iii) physical half-life greater than the time required to prepare material for administration [Iodine 123 (half-life of ~
20 13.2 hours), Iodine 131 (half-life of ~8 days), Gallium 67 (half-life of ~78 hours), and Indium 111 (half-life of ~2.8 days)], (iv) effective half life longer than the examination time, suitable chemical form and reactivity, low toxicity, and stability or near stability of annexin labeled with that radioisotope.

25 *Coupling of Annexin to Contrast Agents*

Coupling of annexin to contrast agents may be performed using any of the art known techniques, *e.g.*, chemical chelation techniques.

Coupling of annexin to a metal oxide may be performed as described in *Chelating Agents and Metal Chelates*, Dwyer & Mellor, Academic Press (1964),
30 Chapter 7 and U.S. Patent No. 5,443,816, the contents of each of which are incorporated herein by reference. Ionic forms of the elements iron, cobalt, nickel, copper, zinc, arsenic, selenium, technetium, ruthenium, palladium, silver, cadmium, indium, antimony, rhenium, osmium, iridium, platinum, gold, mercury, thallium, lead, bismuth, polonium and astatine may be used.

For example, annexin may be incubated with a first reducing agent, the period of incubation being sufficient to reduce available disulfide bonds to thiolate groups while preventing excessive fragmentation of the annexin; the first reducing agent may then be substantially removed from the thiolate-containing annexin; a source of Sn (II) agent
5 may then be added to the thiolate-containing annexin in a sufficient amount to form Sn (II)-containing and sulfur-containing complexes; and the Sn (II)-containing and sulfur-containing complexes may be labeled by adding the metal oxide, whereby the metal oxide displaces the Sn (II) agent and the metal oxide and thiolate-containing annexin form a complex. The order of the foregoing steps may be altered. For example, it is
10 possible, and in some cases advantageous, to add the Sn (II) to form Sn (II)-containing and sulfur-containing complexes prior to removing excess reducing agent from the thiolate-containing annexin. In this way, oxidation of thiolate groups or reformation of disulfide bonds and other cross-linkages can be minimized.

A compound of the invention may be created by associating annexin with
15 biodegradable superparamagnetic metal oxides such as iron oxide. Annexin associated with superparamagnetic or paramagnetic contrast agents provides the advantage of directing the magnetic resonance contrast agent to those cells which are apoptotic or necrotic. A compound prepared from annexin and biodegradable superparamagnetic iron oxide, for example, binds to hepatocytes which are rendered apoptotic by
20 treatment with fas. A magnetic resonance experiment or imaging procedure carried out after administration to a subject of the compounds of the invention can, thus, provide a method for obtaining an enhanced magnetic resonance image, as well as valuable information regarding the distribution of damaged cells in the organism.

The use of magnetic particles for the attachment of biomolecules has been
25 described by Molday (U.S. Patent No. 4,452,773, the entire contents of which are incorporated herein by reference). Briefly, a dextran coated magnetic particle is formed and then treated with periodate to produce aldehyde groups. The aldehydes react with amino groups on a biological molecule, to form a Schiff base. The Schiff base may be stabilized by treatment with a reducing agent like sodium borohydride. After treatment
30 with a reducing agent a methylene amino linker connects the biomolecule to the nanoparticle.

Other methods of attaching biomolecules to nanoparticles, which use the reactivity of the aldehyde group, may also be used, including the methods of Rembaum and Owen (see Table I).

The development of amine functionalized crosslinked iron oxide nanoparticle is another method of synthesizing magnetic particle-biomolecule conjugates that may be used to attach annexins to a metal oxide particle. Amino-CLIO is prepared by first synthesizing a dextran coated magnetic nanoparticle, followed by crosslinking the dextran with epichlorohydrin. Amine groups are incorporated by reacting the dextran with ammonia.

Table I (below) summarizes the types of magnetic particles that may be used for the attachment of Annexins, *e.g.*, Annexin V.

Table I: Magnetic Particles That Can Be Used for the Attachment of Annexins			
Particle Size	Attachment Chem/ Biomolecule Attached	Polymer	Reference
<100 nm	Periodate/antibody	Dextran	Abts (1989) J. Immunol Methods 125,19.
10-70 nm/dextran	Periodate/antibody	Dextran	U.S. Patent No. 4,452,773 (Molday); Molday, (1982) J. Immunol. Methods 52, 353.
10-200 nm/albumin	SPDP/antibody	BSA	U.S. Patent No. 4,795,698 (Owen)
10-50 nm	Periodate/Synaptotagmin 1	Carboxy Dextran	Zhoe (2001) Nat. Med. 7, 1241.
40 nm	SPDP.Oligonucleotides and Peptides	Crosslinked Dextran	Josephson (1999) Bioconjug. Chem. 10, 186.
10-200 nm	Aldehydes/Enzymes, Biomolecules	Polyglutaraldehyde Polymer	U.S. Patent No. 4,438,239 (Rembaum) U.S. Patent No. 4,369,226.
10-100 nm	Periodate/Antibody	Dextran	U.S. Patent No. 5,492,814 (Weissleder)

10

The conjugation of annexins to magnetic molecules yields materials that can be used in a variety of fields such as magnetic affinity chromatography, magnetic cell sorting, magnetic immunoassay and as MR imaging contrast agents. The requirements of the particle vary greatly with the intended application. For imaging applications, the magnetic particle must have a series of properties including:

15

(1) Size and size homogeneity. Magnetic particles are preferably smaller than the size of red blood cells (about 10 microns) to avoid clogging capillary beds. To achieve

efficient targeting to a target cell or organ after injection, they must preferably be in the nanoparticle size range (1-500 nm). Larger particles are rapidly withdrawn from the vascular compartment by the phagocytic cells of the reticuloendothelial system, limiting their ability to react with a limited number of sites on the desired target. Magnetic
5 particles preferably have a narrow size distribution, *i.e.*, cannot have a small percentage of large particles which can occlude capillaries.

(2) Biodegradability. To be useful as a clinical diagnostic tool, magnetic particles must preferably be broken down and excreted or broken down and utilized by the body.

Materials like polystyrene, while useful in the synthesis of magnetic particles for cell
10 sorting, cannot be used in parenteral, clinical applications. The most common type of particle used for imaging applications are polymer coated iron oxides, with dextran or modified dextran being most often employed (Anzai (1994) Radiology 192, 709-15; Reimer (1995) Radiology 195, 489; Stark (1988) Radiology 168, 297).

(3) Safety. The magnetic particles must be non-toxic. Typically, the safety factor (the
15 dose used for imaging divided by the dose killing 50% of a group of animals) is greater than 100 and preferably greater than 1000. Toxicity includes not only the generation of reversible or irreversible tissue damage, but also the induction of transient but annoying physiological reactions in selected subjects (such as humans) taking the preparation. These include fever, urticaria, mild pain, vomiting, and the like. To be useful as a clinical
20 diagnostic agent, such as an MR imaging agent, the magnetic particle must preferably produce no discernable physiological response, except for the desired diagnostic information, in individuals taking preparation.

(4) Stability. To be used as a parenteral agent, the particle must preferably maintain its size distribution during a storage period, which, for practical commercial reasons, is
25 typically longer than 6 months and preferably as long as two years. Instability, evident as the growth in the number of large particles in the preparation, can result in particle induced toxicity, and the abrupt end to the commercial use of the product.

A wide variety of conjugating strategies have been employed to couple proteins to each other and can be adapted to couple Annexins, *e.g.*, Annexin V to magnetic
30 particles, as would be obvious to one skilled in the art. Many of these reagents consist of an N-hydroxysuccinimide ester, which reacts with an amine, and a second moiety that reacts with a sulfhydryl group. A wide selection of bifunctional conjugating reagents, such as SPDP, SMCC, SATA and SIAA are available from Pierce Chemical Company. Detailed procedures for their use are available from the Pierce Chemical web site (see
35 <http://www.piercenet.com>).

Coupling of Annexin to Optically Active Molecules

Coupling of annexin to optically active molecules may be performed using any of the art known techniques, *e.g.*, those described in U.S. Patent No. 5,312,922; U.S. Patent No. 5,928,627; U.S. Patent No. 6,096,289; Weir, ed., Handbook of Experimental
5 Immunology, Vol. 1, Chapter 28, pp. 28.1-28.21, Oxford, Blackwell Scientific, 1986, the entire contents of each of which are incorporated herein by reference.

Administration of the Annexin Containing Compounds of the Invention

The annexin containing compounds of the present invention may be administered to a subject using standard protocols, such as protocols for the
10 administration of radiolabeled compounds.

The compositions of the invention may be administered to a subject in an amount effective, at dosages and for periods of time necessary, to achieve the desired result. An effective amount of the compositions of the invention may vary according to factors such as the tumor stage, age, and weight of the subject, and the ability of the
15 composition to elicit a desired response in the subject. An effective amount is also one in which any toxic or detrimental effects (*e.g.*, side effects) of the compositions are outweighed by the therapeutically or diagnostically beneficial effects. The compositions of the invention may be administered at a concentration of 10-1000 µg protein/kg, 10-900 µg protein/kg, 10-800 µg protein/kg, 10-700 µg protein/kg, 10-600 µg protein/kg,
20 10-500 µg protein/kg, 10-400 µg protein/kg, 10-300 µg protein/kg, 10-200 µg protein/kg, or 10-100 µg protein/kg.

Annexin V begins to have pharmacological effects (anti-coagulant effects) at doses greater than about 300 µg/kg. Accordingly, the diagnostic methods of the present invention (which seek to avoid pharmacological effects of the labeled annexin) are
25 preferably practiced at doses lower than 300 µg/kg, typically less than about 50 µg/kg. Such tracer doses (*e.g.*, 10 µg/kg to 50 µg/kg) have no reported pharmacologic or toxic side effects in animal or human subjects.

The compounds of the invention are typically suspended in a suitable delivery vehicle, such as sterile saline. The vehicle may also contain stabilizing agents, carriers,
30 excipients, stabilizers, emulsifiers, and the like, as is recognized in the art.

The compounds of the invention may be administered to a subject by any suitable route for administration. A preferred method of administration is intravenous (i.v.) injection. It is particularly suitable for imaging of well-vascularized internal organs, such as the heart, liver, spleen, and the like. Methods for i.v. injection of, *e.g.*,
35 radiopharmaceuticals are known. For example, it is recognized that a radiolabeled pharmaceutical is typically administered as a bolus injection using either the

Oldendorf/Tourniquet method or the intravenous push method (see, *e.g.*, Mettler and Guierbteau, (1985) *Essentials Of Nuclear Medicine Imaging*, Second Edition, W.B. Saunders Company, Philadelphia, PA).

For imaging the brain, the compositions of the invention can be administered
5 intrathecally. Intrathecal administration delivers a compound directly to the sub-arachnoid space containing cerebral spinal fluid (CSF). Delivery to spinal cord regions can also be accomplished by epidural injection to a region of the spinal cord exterior to the arachnoid membrane.

For bronchoscopy applications, the annexin compounds of the present invention
10 may be administered by inhalation. For example, the annexin compounds may be delivered in the form of an aerosol spray from a pressured container or dispenser which contains a suitable propellant, *e.g.*, a gas such as carbon dioxide, or a nebulizer.

Other modes of administration include intraperitoneal (*e.g.*, for patients on kidney dialysis), and intrapleural administration. For specific applications, the invention
15 contemplates additional modes of delivery, including intramuscular injection, subcutaneous, intralymphatic, insufflation, and oral, intravaginal and/or rectal administration.

Localization of the Annexin Containing Compounds of the Invention

20 After the compounds of the invention are administered, they are allowed to localize to the target tissue or organ. Localization in this context refers to a condition when either an equilibrium or a pseudo-steady state relationship between bound, "localized", and unbound, "free" compound within a subject has been achieved. The amount of time required for such localization is typically on the order of minutes to tens
25 of minutes and may be estimated by the serum half-life of the compound. The localization time also depends on the accessibility of the target tissue to the compound. This, in turn, depends on the mode of administration, as is recognized in the art.

Imaging is preferably initiated after most of the compound has localized to its target(s). For intravenously administered Tc99m-labeled annexin V, this occurs after
30 several half-lives. A duration of about 10 half-lives (about 30-240 min in the case of annexin/Tc99m conjugates) is considered to be ample time to achieve essentially complete localization. One of skill in the art will appreciate, however, that it may be desirable to perform the imaging at times less than or greater than the ~ 10 half-life timepoint described above. For example, in imaging cell death due to blood vessel
35 injury, the accessibility of the target tissue is very high, such that a strong signal can be

obtained from the target site in only a few minutes, especially if a low dose of labeled annexin is administered gradually to minimize signal from circulating label.

In all of the above cases, a reasonable estimate of the time to achieve localization may be made by one skilled in the art. Furthermore, the state of localization as a
5 function of time may be followed by imaging the gamma ray signal from the labeled annexin according to the methods of the invention.

Applications

Major uses for the annexin containing compounds of the invention include the
10 detection of inappropriate apoptosis in disease states where it should not occur, *e.g.*, immune disorders such as Lupus, transplant rejection, or in cells subject to severe ischemia; and the detection of insufficient apoptosis when it should occur, *e.g.*, tumors or cells infected with a virus.

The annexin containing compounds of the invention may be employed in a
15 variety of clinical settings in which apoptotic and/or necrotic cell death need to be monitored, such as, without limitation, organ and bone marrow transplant rejection or injury, infectious and non-infectious inflammatory diseases, autoimmune disease, cerebral and myocardial infarction and ischemia, cardiomyopathies, atherosclerative disease, neural and neuromuscular degenerative diseases, sickle cell disease,
20 β -thalassemia, cancer therapy, AIDS, myelodysplastic syndromes, and toxin-induced liver disease, and the like. The annexin containing compounds of the invention may also be useful as a clinical research tool to study the normal immune system, embryological development, and immune tolerance and allergy.

The compounds of the invention can be used, for example, to image and quantify
25 apoptotic cell death in normal and malignant tissues undergoing treatment. Monitoring apoptosis with serial imaging studies using these compounds can be used for the rapid testing and development of new drugs and therapies in a variety of diseases. In addition, the methods may be used to monitor the progress of treatment, monitor the progress of disease, or both. Further, they may be used to aid in early detection of certain diseases.

An advantage of the above method is that, by imaging at selected intervals, the
30 method can be used to track changes in the intensity of the emission from the subject over time, reflecting changes in the number of cells undergoing cell death. Such an approach may also be used to track changes in the localization of the compounds of the invention in the subject over time, reflecting changes in the distribution of cells
35 undergoing cell death.

The compositions and methods of the present invention may also be used in the diagnosis and/or treatment of subjects suffering from an eye disease, such as, for example, retinal disease or glaucoma.

The photodynamic therapy (PDT) methods disclosed herein are particularly
5 useful for treating a range of diseases characterized by rapidly growing tissue, including the formation of abnormal blood vessels, such as cancer and age-related macular degeneration (AMD). The type of light source used in PDT varies according to the condition treated. For example, for ophthalmology applications, diode laser light may be shone through the slit lamp of a microscope into a subject's eye. For cancer/ internal
10 diseases, fiber optics may be used to deliver light to the internal cavities like the lung, the gastro-intestinal tract and esophagus and light-emitting diodes (LED) may be used for skin cancer.

In summary, the compositions and methods of the present invention provide a number of clinical and diagnostic benefits. For example, using the methods of the
15 invention, the response of individual patients to established therapeutic anti-cancer regimens may be efficiently and timely evaluated; the anti-neoplastic activity of new anti-cancer drugs may be evaluated; the optimal dose and dosing schedules for new anti-cancer drugs may be identified; and the optimal dose and dosing schedules for existing anti-cancer drugs and drug combinations may be identified. In addition, using the
20 methods of the invention, cancer patients in clinical trials may be categorized efficiently into responders and non-responders to therapeutic regimens.

The methods of the invention provide, among other things, a non-invasive technique for evaluating the early response of individual patient tumors to chemotherapy. This facilitates the selection of effective treatment by allowing rapid
25 identification of ineffective treatments whose side effects might not be balanced by expected benefits.

The invention is further illustrated by the following examples, which should not
30 be construed as further limiting. The contents of all references, pending patent applications and published patents, cited throughout this application, as well as the Figures are hereby expressly incorporated by reference.

EXAMPLES

EXAMPLE 1: Attachment Of Annexin V To Dextran Coated Magnetic Iron Oxides Through The Use Of Periodate

5 Periodate treatment of the dextran coated magnetic particle produces an aldehyde, which forms a Schiff base with the amines of the Annexin V. The complex is stabilized by treatment with sodium borohydride.

 A dextran coated superparamagnetic iron oxide nanoparticle was synthesized according to the methods of Molday (1982) *J. Immunol. Methods* 52, 353. Iron oxide
10 (10 mg Fe in about 1 mL of water) and purified Annexin V were dialyzed against sodium acetate (0.01M, pH 6). Annexin V was purified by the method of Wood (1996) *Blood* 88, 1873. The amount of Annexin V can be varied from 1 to about 50 mg, preferably 5-10 mg of protein. At lower amounts the ratio of protein to iron on the resulting magnetic nanoparticle will be lower, but the offered protein will couple more
15 efficiently. At higher amounts of protein, the ratio of protein to iron on the resulting nanoparticle will be higher, but the percent of protein coupled will be lower.

 Freshly made sodium periodate (50 mg/mL, 0.2 mL) was added to the iron oxide. The mixture was then incubated for 30 minutes at room temperature in the dark, and dialyzed against 0.15 M NaCl. The oxidized magnetic iron oxide was then mixed
20 with the Annexin V and the pH adjusted by the addition of 100 µl of 0.2 M sodium bicarbonate, pH 9.5. The mixture was incubated for 3 hours with stirring. Freshly made sodium cyanoborohydride was then added (25 mg/mL, 0.2 mL) and the mixture was incubated for 6 hours at room temperature. The Annexin V-magnetic nanoparticle can be separated from the unreacted Annexin by a variety sized based separation methods.
25 These include gel filtration, ultrafiltration or magnetic separation.

EXAMPLE 2: Attachment Of Annexin V With A Sulfhydryl Group To Amino CLIO

30 The amino-CLIO nanoparticle was made as described in Josephson (1999) *Bioconjug. Chem.* 10, 186. Annexin V with a sulfhydryl group added through mutagenesis (Tait (2000) *Bioconjug Chem* 11, 918) was employed. To 1.2 mL of amino-CLIO in (30 mg Fe) was added 1.2 mL of 0.1 M phosphate buffer, pH 7.4, and 2
35 mL of N-succinimidyl 3-(2-pyridyldithio)propionate (SPDP, 25 mM) (Molecular Bio-sciences, Boulder, CO) in DMSO. The mixture was allowed to stand for 60 minutes at

room temperature. Low molecular impurities were removed by PD-10 columns (Sigma Chemical, St- Louis, MO) equilibrated with 0.01 M Tris and 0.02 M citrate, pH 7.4 buffer.

Between 2 and 50 mg of Annexin V was subsequently added to 10 mg Fe of the SPDP activated nanoparticle at room temperature and the mixture was allowed to stand overnight. The Annexin V-magnetic nanoparticle can be separated from the unreacted annexin by a variety sized based separation methods.

EXAMPLE 3: Reaction of Annexin V To Add A Sulfhydryl Group, Followed By Reaction With Amino CLIO

A sulfhydryl group was added to the annexin (obtained as in Example 1) by use of the reagent SATA following the manufacturers instructions, Pierce Chemical Company. Amino-CLIO was reacted with SPDP as in Example 2 and then reacted with the SAT A reacted annexin.

EXAMPLE 4: Attachment of Annexin V to a BSA Coated Magnetic Particle

BSA coated magnetic particles were made as described in U.S. Patent No. 4,795,698. Some of the amine groups of the BSA coating of the magnetic particle are converted to sulfydryl groups by use of the reagent SPDP (see Example 2). SPDP or SATA can then be used to add one or more sulfydryl groups on Annexin V. After treatment of the Annexin V with DTT, to expose a sulfhydryl group, the protein is reacted with the magnetic particle.

Equivalents

Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents to the specific embodiments of the invention described herein. Such equivalents are intended to be encompassed by the following

5 claims.

What is claimed is:

1. A magnetic resonance imaging composition comprising an annexin coupled to a contrast agent.
- 5 2. The composition of claim 1, wherein the contrast agent is a paramagnetic agent.
3. The composition of claim 2, wherein the contrast agent is a gadolinium-chelating group complex.
- 10 4. The composition of claim 3, wherein the contrast agent is a gadolinium-diethylenetriamine penta-acetic acid.
- 15 5. The composition of claim 1, wherein the contrast agent is a superparamagnetic agent.
6. The composition of claim 5, wherein the contrast agent is a metal oxide.
- 20 7. The composition of claim 6, wherein the contrast agent is an iron oxide.
8. The composition of claim 6, wherein the contrast agent is a polymer coated metal oxide.
- 25 9. The composition of claim 8, wherein the contrast agent is a dextran coated iron oxide.
10. The composition of claim 1, wherein the annexin is annexin V.
- 30 11. A composition comprising an annexin coupled to a contrast agent and a radioisotope.
12. The composition of claim 11, wherein the radioisotope is a therapeutic radioisotope.
- 35

13. The composition of claim 12, wherein the therapeutic radioisotope is selected from the group consisting of ^{103}Pd , ^{186}Re , ^{188}Re , ^{90}Y , ^{153}Sm , ^{159}Gd , and ^{166}Ho .

14. The composition of claim 11, wherein the annexin is annexin V.

5

15. The composition of claim 11, wherein the contrast agent is a polymer coated metal oxide.

16. A method for imaging cell death in a mammalian subject *in vivo*,
10 comprising

administering to the subject a magnetic resonance imaging composition comprising annexin coupled to a contrast agent; and

obtaining a magnetic resonance image, wherein said image is a representation of cell death in said mammalian subject.

15

17. The method of claim 16, wherein the magnetic resonance image is obtained between about 5 minutes and about 2 hours after the administration of the magnetic resonance imaging composition.

20

18. The method of claim 16, wherein the magnetic resonance image is obtained between about 12-30 hours after the administration of the magnetic resonance imaging composition.

25

19. The method of claim 16, wherein the cell death is caused by apoptosis.

25

20. The method of claim 16, further comprising obtaining a magnetic resonance image at a plurality of time points, thereby monitoring changes in the number of cells undergoing cell death.

30

21. The method of claim 16, further comprising obtaining a magnetic resonance image at a plurality of time points, thereby monitoring changes in the location of cells undergoing cell death.

35

22. The method of claim 16, wherein the magnetic resonance imaging composition is administered at a concentration of 1-500 μg protein/kg.

23. The method of claim 16, wherein the magnetic resonance imaging composition is administered at a concentration of 1-400 µg protein/kg.

5 24. The method of claim 16, wherein the magnetic resonance imaging composition is administered at a concentration of 1-200 µg protein/kg.

25. The method of claim 16, wherein the magnetic resonance imaging composition is administered intravenously.

10

26. The method of claim 16, wherein the magnetic resonance imaging composition is administered via a method selected from the group consisting of intraperitoneally, intrathecally, intrapleurally, intralymphatically and intramuscularly.

15 27. The method of claim 16, wherein cell death is imaged in an organ of a subject or a portion thereof.

28. The method of claim 16, wherein cell death is imaged in the brain of a subject or a portion thereof.

20

29. The method of claim 16, wherein cell death is imaged in the heart of a subject or a portion thereof.

25 30. The method of claim 16, wherein cell death is imaged in the liver of a subject or a portion thereof.

31. An optical imaging composition comprising an annexin coupled to an optically active molecule.

30 32. The composition of claim 31, wherein the optically active molecule is a fluorescent dye.

33. The composition of claim 32, wherein the fluorescent dye is Fluorescein.

35 34. The composition of claim 31, wherein the optically active molecule is a luminescent molecule.

35. The composition of claim 34, wherein the luminescent molecule is luminol.

5 36. The composition of claim 31, wherein the optically active molecule is a bioluminescent molecule.

37. The composition of claim 36, wherein the bioluminescent molecule is selected from the group consisting of luciferase, luciferin, and aequorin.

10

38. The composition of claim 31, wherein the annexin is annexin V.

39. A method for imaging cell death in a mammalian subject *in vivo*, comprising

15

administering to the subject an optical imaging composition comprising annexin coupled to an optically active molecule;

illuminating the subject with a light source; and

visually monitoring the presence of the optical imaging composition in the subject, thereby obtaining an image, wherein said image is a representation of cell death
20 in said mammalian subject.

40. The method of claim 39, wherein the image is obtained between about 5 minutes and about 2 hours after the administration of the optical imaging composition.

25

41. The method of claim 39, wherein the image is obtained between about 12-30 hours after the administration of the optical imaging composition.

42. The method of claim 39, wherein the cell death is caused by apoptosis.

30

43. The method of claim 39, further comprising obtaining an image at a plurality of time points, thereby monitoring changes in the number of cells undergoing cell death.

44. The method of claim 39, further comprising obtaining an image at a plurality of time points, thereby monitoring changes in the location of cells undergoing cell death.

5 45. The method of claim 39, wherein the optical imaging composition is administered at a concentration of 1-500 µg protein/kg.

46. The method of claim 39, wherein the optical imaging composition is administered at a concentration of 1-400 µg protein/kg.

10

47. The method of claim 39, wherein the optical imaging composition is administered at a concentration of 1-200 µg protein/kg.

15 48. The method of claim 39, wherein the magnetic resonance imaging composition is administered via a method selected from the group consisting of intraperitoneally, intrathecally, intrapleurally, intralymphatically and intramuscularly.

49. The method of claim 39, wherein cell death is imaged in an organ of a subject or a portion thereof.

20

50. The method of claim 39, wherein cell death is imaged in the head of a subject or a portion thereof.

25 51. The method of claim 39, wherein cell death is imaged in the heart of a subject or a portion thereof.

52. The method of claim 39, wherein cell death is imaged in the liver of a subject or a portion thereof.

30 53. The method of claim 39, wherein cell death is imaged in the eye of a subject or a portion thereof.

54. A method of tumor radiotherapy, comprising
administering to a subject bearing a tumor an optical imaging composition
35 comprising annexin coupled to an optically active molecule; and

illuminating the subject with a light source in the presence of oxygen, thereby creating a toxic form of oxygen capable of destroying the tumor.

55. The method of claim 54, wherein the optically active molecule is selected
5 from the group consisting of PHOTOFRIN®, Lutrin, ANTRIN®, FOSCAN®,
aminolevulinic acid, aluminum (III) phthalocyanine tetrasulfonate, Hypericin,
verteporfin, and methylene blue dye.

56. The method of claim 54, wherein the toxic form of oxygen is singlet
10 oxygen.

57. The method of claim 54, wherein the tumor is selected from the group
consisting of a brain tumor, a head tumor, a neck tumor, a breast tumor, an esophagus
tumor, a lung tumor, a pleural cavity tumor, an ovary tumor, an abdominal cavity tumor,
15 a bladder tumor, a prostate tumor, a cervix tumor, and a skin tumor.

Figure 1





Figure 2



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
17 October 2002 (17.10.2002)

PCT

(10) International Publication Number
WO 02/080754 A3

(51) International Patent Classification⁷: **A61K 51/00**,
49/00, A61B 5/055

(21) International Application Number: PCT/US02/10238

(22) International Filing Date: 3 April 2002 (03.04.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/281,277 3 April 2001 (03.04.2001) US

(71) Applicant (*for all designated States except US*): **THE-
SEUS IMAGING CORPORATION** [US/US]; 101 Arch
Street, Suite 1762, Boston, MA 02110 (US).

(72) Inventor; and

(75) Inventor/Applicant (*for US only*): **GREEN, Allan, M.**
[US/US]; 19 Francis Avenue, Cambridge, MA 02138 (US).

(74) Agents: **DECONTI, Giulio, A. Jr.** et al.; Lahive & Cock-
field, LLP, 28 State Street, Boston, MA 02109 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,

CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG,
SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ,
VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR,
GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent
(BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR,
NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments

(88) Date of publication of the international search report:
19 December 2002

*For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*

(54) Title: METHODS FOR USING ANNEXIN FOR DETECTING CELL DEATH IN VIVO AND TREATING ASSOCIATED
CONDITIONS

(57) Abstract: The present invention provides methods and compositions for imaging cell death *in vivo*, as well as methods and
compositions for tumor radiotherapy.



WO 02/080754 A3

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US02/10238

A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : A61K 51/00, 49/00; A61B 5/055

US CL : 424/1.69, 9.34, 9.36, 9.6

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 424/1.69, 9.34, 9.36, 9.6

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
Please See Continuation Sheet

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X --- Y	US 5,627,036 A (REUTELINGSPERGER) 06 May 1997, see abstract and columns 5-6.	1-7, 10-12, 14 and 31-38 -----
Y	US 6,197,278 B1 (BLANKENBERG et al) 06 March 2001, see columns 3-4 and column 11, lines 54+.	8, 9, 14-30 and 39-57 11, 12, 14-30 and 39-53
X --- Y	US 6,171,577 B1 (KASINA et al) 09 January 2001, see abstract, column 6, lines 18+ and columns 19-20.	11-14 ----- 1-57
Y	US 5,344,639 A (CHIN et al) 06 September 1994, column 2, lines 5-9.	8, 9 and 15
Y	US 5,976,535 A (FRITZBERG et al) 02 November 1999, see column 8, lines 39-46 and column 22, lines 50-67.	54-57

☐ Further documents are listed in the continuation of Box C.

☐ See patent family annex.

* Special categories of cited documents:		"T"	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"A"	document defining the general state of the art which is not considered to be of particular relevance	"X"	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"E"	earlier application or patent published on or after the international filing date	"Y"	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"L"	document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"&"	document member of the same patent family
"O"	document referring to an oral disclosure, use, exhibition or other means		
"P"	document published prior to the international filing date but later than the priority date claimed		

Date of the actual completion of the international search

12 July 2002 (12.07.2002)

Date of mailing of the international search report

18 OCT 2002

Name and mailing address of the ISA/US

Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703)305-3230

Authorized officer

Michael G. Hartley

Telephone No. (703) 308-1235

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US02/10238

Continuation of B. FIELDS SEARCHED Item 3:

APS

search terms: annexin, imaging, cell death, apoptosis, MRI, radioimaging, fluorescent imaging.